

List of Odonates from Mata do Baú, Barroso, Minas Gerais, Brazil

Marcos Magalhães de Souza^{1*}, Brígida Souza², Matheus Carvalho Soares de Aguiar Pereira³ and Angelo Barbosa Monteiro Machado⁴

1 Instituto Federal de Educação e Tecnologia do Sul de Minas, Campus Inconfidentes. CEP 37576-000. Inconfidentes, MG, Brazil.

2 Universidade Federal de Lavras, Departamento de Entomologia, Caixa Postal 3037. CEP 37200-000. Lavras, MG, Brazil.

3 Universidade Federal de Lavras, Departamento de Biologia, Caixa Postal 3037. CEP 37200-000. Lavras, MG, Brazil.

4 Universidade Federal de Minas Gerais, Instituto de Ciências Biológicas, Departamento de Zoologia, Caixa Postal 486. CEP 31270-901. Belo Horizonte, MG, Brazil.

* Corresponding author: E-mail: magalhaescajubi@bol.com.br

ABSTRACT: A survey of odonatofauna was carried out in Mata do Baú, a predominantly forested area in Barroso, Minas Gerais, regarded as a priority area for conservation and scientific investigation, as published by Biodiversitas, a Brazilian nongovernmental institution. Sample collection was conducted over twenty-six days from November 2009 to February 2011. Fifty-seven species of Odonata were collected and grouped into 30 genera and nine families. The dominant families were Libellulidae (46.5%), Coenagrionidae (20.6%), and Megapodagrionidae (10.3%). A specimen of *Heteragrion obsoletum* (Selys, 1886) was collected, which to-date is known by a single specimen collected in 1880 and red-listed as endangered. Special attention was given to the presence of five species of the genus *Heteragrion*, strictly limited to lotic forest streams, with two new species. This genus is especially sensitive to environmental deterioration, indicating that the forest stream's environmental conditions are beneficial to the area and create a baseline for future monitoring of similar environments.

INTRODUCTION

Taxonomic studies of regional and state fauna are considered important, not only for zoogeographic purposes, but also for defining conservation strategies and supporting the preparation of red lists. With regard to Brazilian odonates, state lists exist only for São Paulo (Costa *et al.* 2000) and Espírito Santo (Costa and Oldrini 2005). In Minas Gerais published, regional lists are scanty and exist only for Poços de Caldas (Santos 1966). Herein, we present a commented list of odonates collected in Mata do Baú, the primary forested area of Barroso, Minas Gerais. This has been regarded as a priority area for conservation and scientific research in Minas Gerais (Drummond *et al.* 2005).

MATERIAL AND METHODS

The study was carried out in Mata do Baú, an environment characterized by the presence of semi-deciduous Montane Forest, gallery forests along the Rio das Mortes river, and savannah (Menini-Neto *et al.* 2004). Comprising an area of 400 ha, Mata do Baú is the largest fragment of native forest in Barroso, situated in the Campos-das-Vertentes region in center-south Minas Gerais (21°11'13" S, 43°58'34" W) (Figure 1). The climate is Cwb mesotermic according to Köppen, characterized by humid summers and dry winters. The mean annual temperature is 18°C with a maximum of 24.4°C and a minimum of 13.8°C. The mean annual rainfall is 1400 mm and the altitude varies between 900 and 1200 m. Collections were made with entomologic nets in the different physiognomic regions, totaling 26 days from November 2009 to February 2011, over sixteen months of study. Specimens were killed with ethyl acetate. The authorization process number issued by IBAMA, through the SISBIO system is 21090.

Specimens were deposited in the entomological collection of the Universidade Federal de Minas Gerais.

Species richness was determined from the absolute numbers of collected specimens, by Jackknife resampling, using the Estimates Software (Colwell 2009) with 1000 resamplings. Collection efficiency and the number of sampled species were based on the richness percentages estimated by the mean of three nonparametric estimators: Ace 1, Jack 1, and Chao 1 with the use of estimates. The total number of expected species was compared using the species accumulation curve (Cowell 2009), with a confidence degree of 95%, to estimate the actual number of species in the studied area.




FIGURE 1. Geographic localition of Barroso, state of Minas Gerais, Brazil.

RESULTS AND DISCUSSION

The list Odonata recorded from Mata do Baú is present as follows:

ZYGOPTERA**CALOPTERYGIDAE**

Hetaerina longipes Hagen and Selys, 1853 - I-OD-2012-0001

Hetaerina rosea Selys, 1853 - I-OD-2012-0002

Hetaerina simplex Selys, 1853 - I-OD-2012-0003

Mnesarete pudica phryne Costa, 1986 - I-OD-2012-0004

COENAGRIONIDAE

Acanthagrion gracile Rambur, 1842 - I-OD-2012-0005

Argia croceipennis Selys, 1865 - I-OD-2012-0006

Argia lilacina Selys, 1865 - I-OD-2012-0007

Argia mollis Hagen in Selys, 1865 - I-OD-2012-0008

Argia reclusa Selys, 1865 - I-OD-2012-0009

Argia sp. - I-OD-2012-0010

Oxyagrion basale Selys, 1876 - I-OD-2012-0011

Oxyagrion terminale Selys, 1876 - I-OD-2012-0012

Oxyagrion evanescens Calvert, 1909 - I-OD-2012-0013

Telebasis gigantea Daigle, 2002 - I-OD-2012-0015

Tigriagrion aurantinigrum Calvert, 1909 - I-OD-2012-0016

DICTERIADIDAE

Heliogaris amazona Selys, 1853 - I-OD-2012-0017

LESTIDAE

Lestes auritus Hagen in Selys, 1862 - I-OD-2012-0018

Lestes bipupillatus Calvert, 1909 - I-OD-2012-0019

MEGAPODAGRIONIDAE

Allopodagrion contortum (Hagen in Selys, 1862) - I-OD-2012-0020

Heteragrion aurantiacum Selys, 1862 - I-OD-2012-0021

Heteragrion obsoletum Selys, 1886 - I-OD-2012-0022

Heteragrion tiradentense Machado and Bedé, 2006 - I-OD-2012-0023

Heteragrion n. sp. - I-OD-2012-0024

Heteragrion sp. - I-OD-2012-0025

PROTONEURIDAE

Epileoneura venezuelensis Rácenis, 1955 - I-OD-2012-0026

Forcepsioneura sancta (Hagen in Selys, 1860) - I-OD-2012-0027

ANISOPTERA**AESHNIDAE**

Castoraeschna januararia (Hagen, 1857) - I-OD-2012-0028

Rhionaeschna pauloi Machado, 1984 - I-OD-2012-0029

GOMPHIDAE

Epigomphus paludosus Hagen in Selys, 1854 - I-OD-2012-0030

Phyllocycla argentina (Hagen in Selys, 1856) - I-OD-2012-0031

LIBELLULIDAE

Brachymesia furcata (Hagen, 1861) - I-OD-2012-0032

Brechmorhoga nubecula (Rambur, 1842) - I-OD-2012-0033

Dasythemis mincki mincki (Karsch, 1890) - I-OD-2012-0034

Elasmothemis constricta (Calvert, 1898) - I-OD-2012-0035

Erythrodiplax fusca (Rambur, 1842) - I-OD-2012-0036

Erythrodiplax latimaculata Ris, 1911 - I-OD-2012-0037

Erythrodiplax media Borrer, 1942 - I-OD-2012-0038

Erythrodiplax umbrata (Linnaeus, 1758) - I-OD-2012-0039

Erythrodiplax paraguayensis (Förster, 1904) - I-OD-2012-0040

Erythrodiplax juliana Ris, 1911 - I-OD-2012-0041

Erythrodiplax sp. 1 - I-OD-2012-0042

Erythrodiplax sp. 2 - I-OD-2012-0043

Erythemis peruviana Rambur, 1842 - I-OD-2012-0044

Erythemis vesiculosa (Fabricius, 1775) - I-OD-2012-0045

Macrothemis heteronycha (Calvert, 1909) - I-OD-2012-0046

Macrothemis imitans imitans Karsch, 1890 - I-OD-2012-0047

Macrothemis marmorata Hagen, 1868 - I-OD-2012-0048

Miathyria marcella (Selys, 1857) - I-OD-2012-0049

Micrathyria stawiarskii Santos, 1953 - I-OD-2012-0050

Micrathyria hesperis Ris, 1911 - I-OD-2012-0051

Micrathyria sp. - I-OD-2012-0052

Orthemis discolor (Burmeister, 1839) - I-OD-2012-0053

Pantala flavescens (Fabricius, 1798) - I-OD-2012-0054

Perithemis mooma Kirby, 1889 - I-OD-2012-0055

Perithemis icteroptera (Selys, 1857) - I-OD-2012-0056

Tramea calverti Muttkowski, 1910 - I-OD-2012-0057

Tramea cophysa Hagen, 1817 - I-OD-2012-0058

As shown in the list, 57 species belonging to 30 genera and nine families were collected in Mata do Baú. Considering the small size of the area (400 ha), the number of species is relatively large and reflects the diversity of the area, which contains both lotic and lentic water systems, as well as forested and open environments. The results of a social wasps inventory (Vespidae) also revealed a high number of species, a fact that has been correlated with the area's plant diversity (Souza and Prezoto 2006). Table 1 shows the number of species for each family.

Libellulidae was the family with most species (46.55%) followed by Coenagrionidae (20.60%) and Megapodagrionidae (10.3%), in marked contrast with the fact that in the world and in the Neotropical region, Coenagrionidae is more speciose than Libellulidae and Megapodagrionidae. On the other hand, all collected Libellulidae were common species with wide geographic ranges (although three of them – two *Erythrodiplax* and one *Micrathyria* – all represented by male individuals, not yet been identified), whereas the rarest and most important species, with regard to their conservation status, are Coenagrionidae and Megapodagrionidae.

The common species, with wide geographic ranges, predominated in the areas of open vegetation, whereas the rarest and most important for conservation predominated in the forested area, a fact that corroborates the results obtained for the odonatological fauna of the Maraca Biological Reserve in Roraima (Machado *et al.* 1991). In the Megapodagrionidae, five species of *Heteragrion* were collected, all from the gallery forest lotic streams. The presence of the species of this genus is an unusual finding.

Its species are strictly lotic with most species occurring only in lotic, streams, in areas of dense forest. Adults and larval odonates have been shown to be good indicators of water quality (Schmidt 1985; Brown Jr. 1997; Clausnitzer and Jödicke 2004) and *Heteragrion* has been shown to be especially sensitive to environmental deterioration (Machado 1988).

Thus, the presence of five species of *Heteragrion* in Mata do Baú, with a new species under description, indicates that its water systems are well preserved and present a strong argument for creating a protected area in this area. Moreover, the rich presence of *Heteragrion* is a useful tool for its future monitoring.

Of especial interest was finding a specimen of *Heteragrion obsoletum* Selys, 1886 (Figure 02), collected in a shaded area inside the forest. As reported by Machado (2008), this species was described based on a single specimen collected by the Belgian naturalist Walther de Selys-Lonchamps in December, 1872, in Caxambu, Minas Gerais, as recorded in his diary (Selys 1875). The specimen has been deposited in the Selys collection at the Institute Royal des Sciences Naturelles de Belgique in Brussels. Since then, the species has never again been collected, despite the fact that two collection trips have been made to Caxambu and neighboring Airuoca in an attempt to find it (Machado, personal communication). Its rediscovery at Mata do Baú, situated around 150 km from Caxambu, is especially important because the holotype of the single known specimen disappeared from the Selys collection as reported by the curator Jerome Constant (in litt.). The species is regarded as threatened under the endangered category both on the National and State red lists (Machado 2008). Based on the specimen from Barroso, the species will be redescribed and a neotype designated in a separate paper. Another rare species collected at Mata do Baú is *Heteragrion tiradentense* (Machado and Bedê 2006) (Figure 3) known from a single locality at the Serra de São José Wildlife Refuge in Tiradentes, Minas Gerais.

The finding of *Heliocharis amazona* Selys, 1853 (Dicteriadidae), restricted to streams of the well-preserved stretches of gallery forests along the Rio das Mortes river, suggests that this species may also be a good indicator of the degree of conservation of the streams, as already pointed out by Ferreira-Peruquetti and Fonseca-Gessner (2003) (Figure 4).

Despite the fact that the number of species obtained was relatively high, collection efficiency by means of nonparametric estimators was 61%, indicating that collection effort was not enough to reveal the true richness of odonatofauna at Serra do Baú. Taking our knowledge about the Odonata fauna in environments similar to those of the Mata do Baú into account, we can state that further collection efforts would certainly allow other species to be found, especially in the Rio das Mortes, in which one or two species of *Neoneura*, one species of *Elasmothermis*, and other species of *Gomphidae* possibly occur. One should also consider that, when analyzed separately, the estimator 01 Jack differs from that of the collected species curve (cole) and indicates the stability of the species collected (Figure 5).



FIGURE 2. *Heteragrion obsoletum* Selys, 1886



FIGURE 3. *Heteragrion tiradentense* (Machado and Bedê 2006)



FIGURE 4. *Heliocharis amazona* Selys, 1853




FIGURE 5. Number of species of Odonata collected (sobs) on 26 sampling days in relation to the number of species expected by Jackknife 01 estimator in Mata do Baú, Barroso, state of Minas Gerais, Brazil.

TABLE 1. Number of Odonata species per family collected in Mata do Baú.

FAMILY	No.	%
Zygoptera		
Calopterygidae	4	6.99
Coenagrionidae	11	20.60
Dictyrididae	1	1.91
Megapodagrionidae	6	10.30
Protoneuridae	2	3.44
Lestidae	2	3.44
Anisoptera		
Aeschnidae	2	3.44
Gomphidae	2	3.44
Libellulidae	27	46.55
TOTAL	58	100%

ACKNOWLEDGMENTS: We thank FAPEMIG for the financial support to the Project (Programa Biota Minas, edital 14/2009); IBAMA for the collection permit; Melo family, owners of "Mata do Baú", for permitting the studies in this area; the staff members of the Project, Marcell Santos Ferreira and Epifânio Porfiro Pires, for helping with the field collections; and Dr. Abner Elpino Campos, Universidade de Concórdia, SC, for help with statistical tests.

LITERATURE CITED

Brown-Jr., K.S. 1997. Insetos como rápidos e sensíveis indicadores de uso sustentável de recursos naturais; p. 143-155. In H.L. Martos and N.B. Maia (ed.). *Indicadores ambientais*. Sorocaba: PUC/SP.

Clausnitzer, V. and R. Jödicke. 2004. Guardians of the watershed. *International Journal of Odonatology* 7(2): 111.

Colwell, R.K. 2009. *EstimateS: statistical estimation of species richness and shared species from samples, version 6.0 b1: user's guide and application*. Storrs: University of Connecticut. Electronic Database accessible at <http://viceroy.eeb.uconn.edu/estimates> 2009. Captured on 12 December 2011.

Costa, J.M., A.B.M. Machado, F.A.A. Lencioni and T.C. Santos. 2000. Diversidade e distribuição dos Odonata (Insecta) no Estado de São Paulo, Brasil: Parte I – Lista das espécies e registros bibliográficos. *Publicações Avulsas do Museu Nacional* 80: 1-27.

Costa, J.M. and B.B. Oldrini. 2005. Diversidade e distribuição dos Odonata (Insecta) no Estado do Espírito Santo, Brasil. *Publicações Avulsas do Museu Nacional* 107: 1–15.

Drummond, G.M., C.S. Martins, A.B.M. Machado, F.A. Sebaio and Y. Antonini. 2005. *Biodiversidade em Minas Gerais. Um Atlas para sua Conservação*. Belo Horizonte: Fundação Biodiversitas. 222 p.

Ferreira-Peruquetti, P. S. and A. A. Fonseca-Gessner. 2003. Comunidade de Odonata (Insecta) em áreas naturais de Cerrado e monocultura no nordeste do Estado de São Paulo, Brasil: relação entre o uso do solo e a riqueza faunística. *Revista Brasileira de Zoologia* 20: 219–224.

Machado, A.B.M. 1988. *Heteragrion petiense* spec. nov. from the state of Minas Gerais, Brazil (Zygoptera: Megapodagrionidae). *Odonatologica* 17: 267-274.

Machado, A.B.M., H.G. Mesquita and P.A.R. Machado. 1991. Contribuição ao conhecimento dos odonatos da Estação Ecológica de Maracá. *Acta Amazônica* 21: 159-173.

Machado, A.B.M. and L.C. Bedê. 2006. *Heteragrion tiradentense* spec. nov. from the state of Minas Gerais, Brazil (Zygoptera: Megapodagrionidae). *Odonatologica* 35: 47-54.

Machado, A.B.M. 2008. *Heteragrion obsoletum* Selys, 1886; p. 459-460. In A.B.M. Machado, G.M. Drummond and A.P. Pageia (ed.). *Livro Vermelho da Fauna Brasileira Ameaçada de Extinção*. Volume I. Belo Horizonte: Fundação Biodiversitas.

Menini-Neto, M.L., L.C.S. Assis and R.C. Forzza. 2004. A família Orchidaceae em um fragmento de floresta estacional semidecidual, no município de Barroso, Minas Gerais, Brasil. *Lundiana* 4: 9-27.

Santos, N.D. 1966. Odonatos da região de Poços de Caldas, Minas Gerais. *Atas da Sociedade de Biologia do Rio de Janeiro* 10: 65-68.

Schmidt, E. 1985. Habitat invitarization characterlization and bioindication by a representative spectrum of Odonata species. *Odonatologica* 14(2): 127-133.

Selys, W. de- 1875. Notes d'un Voyage au Brésil. Bruxelles, Librairie C. Muaquardt 102 pp.

Souza, M.M. and F. Prezoto. 2006. Diversity of social wasps (Hymenoptera, Vespidae) in semideciduous forest and cerrado (savanna) regions in Brazil. *Sociobiology* 47: 135-147.

RECEIVED: September 2012

ACCEPTED: October 2013

PUBLISHED ONLINE: November 2013

EDITORIAL RESPONSIBILITY: Rodolfo Mariano